arXiv:1701.02164v1 [math.RA] 9 Jan 2017

Symmetric square-central elements in products of orthogonal involutions in characteristic two

A.-H. Nokhodkar

Abstract

We obtain some criteria for a symmetric square-central element of a totally decomposable algebra with orthogonal involution in characteristic two, to be contained in an invariant quaternion subalgebra.
Mathematics Subject Classification: 16W10, 16K20, 11E39.

1 Introduction

A classical question concerning central simple algebras is to identify under which conditions a square-central element lies in a quaternion subalgebra. This question is only solved for certain special cases in the literature. Let A be a central simple algebra of exponent two over a field F. In [2, (3.2)] it was shown that if $\operatorname{char} F \neq 2$ and F is of cohomological dimension less than or equal 2, then every square-central element of A is contained in a quaternion subalgebra (see also [4, (4.2)]). In [3, (4.1)], it was shown that if $\operatorname{char} F \neq 2$, then an element $x \in A$ with $x^2 = \lambda^2 \in F^{\times 2}$ lies in a (split) quaternion subalgebra if and only if $\dim_F (x-\lambda)A = \frac{1}{2}\dim_F A$. This result was generalized in [16, (3.2)] to arbitrary characteristics, including $\lambda = 0$. On the other hand, in [19] it was shown that there is an indecomposable algebra of degree 8 and exponent 2, containing a square-central element (see [9, (5.6.10)]). Using similar methods, it was shown in [3] that for $n \geqslant 3$ there exists a tensor product of n quaternion algebras containing a square-central element which does not lie in any quaternion subalgebra.

A similar question for a central simple algebra with involution (A, σ) over F is whether a symmetric or skew-symmetric square-central element of A lies in a σ-invariant quaternion subalgebra. In the case where $\operatorname{char} F \neq 2$, $\deg_F A = 8$, σ has trivial discriminant, the index of A is not 2 and one of the components of the Clifford algebra $C(A, \sigma)$ splits, it was shown in [18, (3.14)] that every skew-symmetric square-central element of A lies in a σ-invariant quaternion subalgebra. In [15], some criteria were obtained for symmetric and skew-symmetric elements whose squares lie in F^2, to be contained in a σ-invariant quaternion subalgebra. Also, a sufficient condition was obtained in [14, (6.3)] for symmetric square-central elements in a totally decomposable algebra with orthogonal involution in characteristic two, to be contained in a stable quaternion subalgebra.

In this work we study the aforementioned question for totally decomposable algebras with orthogonal involution in characteristic two. Let (A, σ) be a totally decomposable algebra with orthogonal involution over a field F of characteristic

two and let $x \in A \setminus F$ be a symmetric element with $\alpha := x^2 \in F$. Since the case where $\alpha \in F^2$ was settled in [15], we assume that $\alpha \in F^\times \setminus F^{\times 2}$. We first study in §3 some properties of *inseparable subalgebras*, introduced in [14]. It is shown in Theorem 3.10 that (A, σ) has a unique inseparable subalgebra if and only if either $\deg_F A \leqslant 4$ or σ is anisotropic. In §4, the notion of the *representation* of an element of F by σ is defined. Theorem 4.10 shows that the set of elements in F represented by σ coincides with the set of elements in F represented by the *Pfister invariant* of (A, σ) (a bilinear Pfister form associated to (A, σ) and defined in [6]). In particular, if $0 \neq x \in A$ is symmetric and square-central, then the element $x^2 \in F$ is represented by the Pfister invariant (see Corollary 4.9).

We then study our main problem in §5 and §6. For the case where σ is anisotropic or A has degree 4, it is shown that every symmetric square-central element of A lies in a σ-invariant quaternion subalgebra (see Theorem 5.1 and Proposition 5.2). However, we will see in Proposition 6.2 that if σ is isotropic, $\deg_F A \geqslant 8$ and $(A, \sigma) \not\simeq (M_{2^n}(F), t)$, there always exists a symmetric square-central of (A, σ) which is not contained in any σ-invariant quaternion subalgebra of A. If A has degree 8 or σ satisfies certain isotropy condition, it is shown in Proposition 5.7 and Theorem 5.10 that a symmetric square-central element of A lies in a σ-invariant quaternion subalgebra if and only if it is contained in an inseparable subalgebra of (A, σ). Finally, in Example 6.3 we shall see that this criterion cannot be applied to arbitrary involutions.

2 Preliminaries

Throughout this paper, F denote a field of characteristic two.

Let A be a central simple algebra over F. An *involution* on A is an anti-automorphism $\sigma : A \to A$ of order two. If $\sigma|_F = \mathrm{id}$, we say that σ is of *the first kind*. The sets of *alternating* and *symmetric* elements of (A, σ) are defined as

$$\mathrm{Sym}(A, \sigma) = \{x \in A \mid \sigma(x) = x\} \quad \text{and} \quad \mathrm{Alt}(A, \sigma) = \{\sigma(x) - x \mid x \in A\}.$$

For a field extension K/F we use the notation $A_K = A \otimes K$, $\sigma_K = \sigma \otimes \mathrm{id}$ and $(A, \sigma)_K = (A_K, \sigma_K)$. An extension K/F is called a *splitting field* of A if A_K splits, i.e., A_K is isomorphic to the matrix algebra $M_n(K)$, where $n = \deg_F A$ is the degree of A over F. If $(V, \mathfrak{b})$ is a symmetric bilinear space over F, the pair $(\mathrm{End}_F(V), \sigma_\mathfrak{b})$ is denoted by $\mathrm{Ad}(\mathfrak{b})$, where $\sigma_\mathfrak{b}$ is the *adjoint involution* of $\mathrm{End}_F(V)$ with respect to $\mathfrak{b}$ (see [10, p. 2]). According to [10, (2.1)], if K is a splitting field of A, then $(A, \sigma)_K$ is adjoint to a symmetric bilinear form. We say that σ is *symplectic*, if this form is alternating. Otherwise, σ is called *orthogonal*. By [10, (2.6)], σ is symplectic if and only $1 \in \mathrm{Alt}(A, \sigma)$. If σ is an orthogonal involution, the *discriminant* of σ is denoted by $\mathrm{disc}\, \sigma$ (see [10, (7.2)]).

Let $(V, \mathfrak{b})$ be a bilinear space over F and let $\alpha \in F$. We say that $\mathfrak{b}$ *represents* α if $\mathfrak{b}(v, v) = \alpha$ for some nonzero vector $v \in V$. The set of elements in F represented by $\mathfrak{b}$ is denoted by $D(\mathfrak{b})$. We also set $Q(\mathfrak{b}) = D(\mathfrak{b}) \cup \{0\}$. Observe that $Q(\mathfrak{b})$ is an F^2-subspace of F. If K/F is a field extension, the scalar extension of $\mathfrak{b}$ to K is denoted by $\mathfrak{b}_K$. For $\alpha_1, \cdots, \alpha_n \in F^\times$, the diagonal bilinear form $\sum_{i=1}^n \alpha_i x_i y_i$ is denoted by $\langle \alpha_1, \cdots, \alpha_n \rangle$. The form $\langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle := \langle 1, \alpha_1 \rangle \otimes \cdots \otimes \langle 1, \alpha_n \rangle$ is called a *bilinear (n-fold) Pfister form*. If $\mathfrak{b}$ is a bilinear Pfister form over F, then there exists a bilinear form $\mathfrak{b}'$, called the *pure subform*

of $\mathfrak{b}$, such that $\mathfrak{b} \simeq \langle 1 \rangle \perp \mathfrak{b}'$. The form $\mathfrak{b}'$ is uniquely determined, up to isometry (see [1, p. 906]).

3 The inseparable subalgebra

Definition 3.1. An involution σ on a central simple algebra A is called *isotropic* if $\sigma(x)x = 0$ for some nonzero element $x \in A$. Otherwise, σ is called *anisotropic*.

The following definition was given in [5, p. 1629].

Definition 3.2. A central simple algebra with involution (A, σ) is called *direct* if $\sigma(x)x \in \mathrm{Alt}(A, \sigma)$ for $x \in A$ implies that $x = 0$.

It is clear that every direct algebra with involution is anisotropic.

Notation 3.3. For an algebra with involution (A, σ) over F we use the following notation:

$$\mathrm{Alt}(A, \sigma)^+ = \{x \in \mathrm{Alt}(A, \sigma) \mid x^2 \in F\},$$
$$\mathrm{Sym}(A, \sigma)^+ = \{x \in \mathrm{Sym}(A, \sigma) \mid x^2 \in F\}.$$

Note that we have $\mathrm{Alt}(A, \sigma)^+ \oplus F \subseteq \mathrm{Sym}(A, \sigma)^+$.

Proposition 3.4. *If (A, σ) is a direct algebra with orthogonal involution over F, then $\mathrm{Sym}(A, \sigma)^+$ is a subfield of A.*

Proof. Let $x, y \in \mathrm{Sym}(A, \sigma)^+$. Set $\alpha = x^2 \in F$ and $\beta = y^2 \in F$. Consider the element $w := xy + yx \in \mathrm{Sym}(A, \sigma)$. We have

$$\begin{aligned}\sigma(w)w = w^2 &= (xy + yx)^2 = xyxy + yxyx + xy^2x + yx^2y \\ &= xyxy + yxyx + \alpha\beta + \alpha\beta = \sigma(yxyx) - yxyx \in \mathrm{Alt}(A, \sigma).\end{aligned}$$

Hence $w = 0$, i.e., $xy = yx$. It follows that $\sigma(xy) = xy$ and $(xy)^2 \in F$, i.e., $xy \in \mathrm{Sym}(A, \sigma)^+$. We also have $(x + y)^2 = \alpha + \beta + xy + yx = \alpha + \beta \in F$, hence $x + y \in \mathrm{Sym}(A, \sigma)^+$. The set $\mathrm{Sym}(A, \sigma)^+$ is therefore a commutative subalgebra of A with $u^2 \in F$ for every $u \in \mathrm{Sym}(A, \sigma)^+$. Since σ is direct, if $0 \neq u \in \mathrm{Sym}(A, \sigma)^+$, then $u^2 \neq 0$, i.e., u is a unit. Hence, $\mathrm{Sym}(A, \sigma)^+$ is a field. □

A central simple algebra with involution is called *totally decomposable* if it decomposes into tensor products of quaternion algebras with involution. Let $(A, \sigma) \simeq \bigotimes_{i=1}^{n} (Q_i, \sigma_i)$ be a totally decomposable algebra of degree 2^n with orthogonal involution over F. By [10, (2.23)] the involution σ_i is necessarily orthogonal for $i = 1, \cdots, n$. Also, according to [14, (4.6)] there exists a 2^n-dimensional subalgebra $S \subseteq \mathrm{Sym}(A, \sigma)^+$, called an *inseparable subalgebra* of (A, σ), satisfying (i) $C_A(S) = S$, where $C_A(S)$ is the centralizer of S in A; and (ii) S is generated, as an F-algebra, by n elements. Furthermore, for every inseparable subalgebra S of (A, σ) we have necessarily $S \subseteq \mathrm{Alt}(A, \sigma)^+ \oplus F$. According to [14, (5.10)], if S_1 and S_2 are two inseparable subalgebras of (A, σ), then $S_1 \simeq S_2$ as F-algebras. Note that if $v_i \in \mathrm{Sym}(Q_i, \sigma_i)^+ \setminus F$ is a unit for $i = 1, \cdots, n$, then $F[v_1, \cdots, v_n]$ is an inseparable subalgebra of (A, σ).

Theorem 3.5. *Let (A,σ) be a totally decomposable algebra with anisotropic orthogonal involution over F and let S be an inseparable subalgebra of (A,σ). Then $S = \mathrm{Alt}(A,\sigma)^+ \oplus F = \mathrm{Sym}(A,\sigma)^+$ is a maximal subfield of A. In particular, the inseparable subalgebra S is uniquely determined.*

Proof. We already know that $S \subseteq \mathrm{Alt}(A,\sigma)^+ \oplus F \subseteq \mathrm{Sym}(A,\sigma)^+$. By [6, (6.1)], (A,σ) is direct. Hence, $\mathrm{Sym}(A,\sigma)^+$ is a field by Proposition 3.4. Since S is maximal commutative, we obtain $\mathrm{Sym}(A,\sigma)^+ = S$. Finally, as $\dim_F S = \deg_F A$, S is a maximal subfield of A. □

The following definition was given in [6].

Definition 3.6. Let $(A,\sigma) \simeq \bigotimes_{i=1}^{n}(Q_i,\sigma_i)$ be a totally decomposable algebra with orthogonal involution over F. The *Pfister invariant* of (A,σ) is defined as $\mathfrak{Pf}(A,\sigma) := \langle\!\langle \alpha_1,\cdots,\alpha_n\rangle\!\rangle$, where $\alpha_i \in F^\times$ is a representative of the class $\mathrm{disc}\,\sigma_i \in F^\times/F^{\times 2}$, $i = 1,\cdots,n$.

According to [6, (7.2)], the isometry class of the Pfister invariant is independent of the decomposition of (A,σ). Moreover, if S is an inseparable subalgebra of (A,σ), then S may be considered as an underlying vector space of $\mathfrak{Pf}(A,\sigma)$ such that $\mathfrak{Pf}(A,\sigma)(x,x) = x^2$ for $x \in S$ (see [14, (5.5)]).

Notation 3.7. For a positive integer n, we denote the bilinear n-fold Pfister form $\langle\!\langle 1,\cdots,1\rangle\!\rangle$ by $\langle\!\langle 1\rangle\!\rangle^n$. We also set $\langle\!\langle 1\rangle\!\rangle^0 = \langle 1\rangle$.

Let $\mathfrak{b}$ be a bilinear Pfister form over F. In view of [1, A.5], one can find a non-negative integer r and an anisotropic bilinear Pfister form $\mathfrak{b}'$ such that $\mathfrak{b} \simeq \langle\!\langle 1\rangle\!\rangle^r \otimes \mathfrak{b}'$. As in [15], we denote the integer r by $\mathfrak{i}(\mathfrak{b})$. If (A,σ) is a totally decomposable F-algebra with orthogonal involution, we simply denote $\mathfrak{i}(\mathfrak{Pf}(A,\sigma))$ by $\mathfrak{i}(A,\sigma)$. By [6, (5.7)], (A,σ) is anisotropic if and only if $\mathfrak{i}(A,\sigma) = 0$. Hence, if σ is isotropic, then the pure subform of $\mathfrak{Pf}(A,\sigma)$ represents 1. In other words, if S is an inseparable subalgebra of (A,σ), then there exists $x \in S \setminus F$ with $x^2 = 1$. Also, if $r = \mathfrak{i}(A,\sigma) > 0$, there exists a totally decomposable algebra with anisotropic orthogonal involution (B,ρ) over F such that $(A,\sigma) \simeq (M_{2^r}(F),t) \otimes (B,\rho)$, where t is the transpose involution (see [15, p. 7]). In particular, if A is of degree 2^n then $\mathfrak{i}(A,\sigma) = n$ if and only if $(A,\sigma) \simeq (M_{2^n}(F),t)$.

We recall that every quaternion algebra Q over F has a *quaternion basis*, i.e., a basis $(1,u,v,w)$ satisfying $u^2+u \in F$, $v^2 \in F^\times$ and $w = uv = vu+v$ (see [10, p. 25]). In this case, Q is denoted by $[\alpha,\beta)_F$, where $\alpha = u^2+u \in F$ and $\beta = v^2 \in F^\times$.

Lemma 3.8. *If (Q,σ) is a quaternion algebra with orthogonal involution over F, then there is a quaternion basis $(1,u,v,w)$ of Q such that $u,v \in \mathrm{Sym}(Q,\sigma)$.*

Proof. Let $v \in \mathrm{Alt}(Q,\sigma)$ be a unit. Since $v \notin F$ and $v^2 \in F^\times$, it is easily seen that v extends to a quaternion basis $(1,u,v,w)$ of Q. By [15, (4.5)], we have $\sigma(u) = u$. □

Lemma 3.9. *Let (A,σ) be a totally decomposable algebra of degree 8 with orthogonal involution over F. If σ is isotropic, then there are two inseparable subalgebras S_1 and S_2 of (A,σ) with $S_1 \neq S_2$.*

Proof. Since $\mathfrak{i}(A,\sigma) > 0$, we may identify $(A,\sigma) = (B,\rho) \otimes (M_2(F),t)$, where $(B,\rho) \simeq (Q_1,\sigma_1) \otimes (Q_2,\sigma_2)$ is a tensor product of two quaternion algebras with orthogonal involution. By Lemma 3.8 there exists a quaternion basis $(1,u_i,v_i,w_i)$ of Q_i over F such that $u_i, v_i \in \mathrm{Sym}(Q_i,\sigma_i)$, $i = 1,2$. Let $v_3 \in \mathrm{Alt}(M_2(F),t)$ be a unit. By scaling we may assume that $v_3^2 = 1$, because $\operatorname{disc} t$ is trivial (see [10, p. 82]). Then $S_1 = F[v_1 \otimes 1, v_2 \otimes 1, 1 \otimes v_3]$ is an inseparable subalgebra of (A,σ) (we have identified Q_1 and Q_2 with subalgebras of B). Let

$$v_1' = u_2v_1 \otimes 1 + (u_2v_1 + v_1) \otimes v_3 \in \mathrm{Sym}(A,\sigma)^+,$$
$$v_2' = u_1v_2 \otimes 1 + (u_1v_2 + v_2) \otimes v_3 \in \mathrm{Sym}(A,\sigma)^+.$$

Computation shows that $S_2 = F[v_1', v_2', 1 \otimes v_3]$ is another inseparable subalgebra of (A,σ). Clearly, we have $S_1 \neq S_2$. □

Theorem 3.10. *A totally decomposable algebra with orthogonal involution (A,σ) over F has a unique inseparable subalgebra if and only if either $\deg_F A \leqslant 4$ or σ is anisotropic.*

Proof. Let S be an inseparable subalgebra of (A,σ). If A is a quaternion algebra, then $S = \mathrm{Alt}(A,\sigma) \oplus F$ by dimension count. If $\deg_F A = 4$, then $S = \mathrm{Alt}(A,\sigma)^+ \oplus F$ by [17, (4.4)]. Also, if σ is anisotropic, then S is uniquely determined by Theorem 3.5. This proves the 'if' implication. To prove the converse, let $\deg_F A = 2^n$. Suppose that σ is isotropic and $\deg_F A \geqslant 8$, i.e., $n \geqslant 3$. As $\mathfrak{i}(A,\sigma) > 0$, we may identify $(A,\sigma) = \bigotimes_{i=1}^{n-1}(Q_i,\sigma_i) \otimes (M_2(F),t)$, where (Q_i,σ_i) is a quaternion algebra with orthogonal involution over F for $i = 1,\cdots,n-1$. By Lemma 3.9 the algebra with involution

$$(Q_{n-2},\sigma_{n-2}) \otimes (Q_{n-1},\sigma_{n-1}) \otimes (M_2(F),t),$$

admits two inseparable subalgebras S_1 and S_2 with $S_1 \neq S_2$. Let S be an inseparable subalgebra of $\bigotimes_{i=1}^{n-3}(Q_i,\sigma_i)$. Then $S \otimes S_1$ and $S \otimes S_2$ are two inseparable subalgebras of (A,σ) with $S \otimes S_1 \neq S \otimes S_2$, proving the result. □

Lemma 3.11. *Let (A,σ) be a totally decomposable algebra of degree 2^n with orthogonal involution over F and let $x \in \mathrm{Sym}(A,\sigma)^+$ be a unit. If S is an inseparable subalgebra of (A,σ), then for every unit $y \in S$, there exists a positive integer k such that $(xy)^k \in \mathrm{Sym}(A,\sigma)^+$. In addition, for such an integer k we have $(xy)^k x = x(xy)^k$.*

Proof. Since x and y are units, the element $(xy)^r$ is a unit for every integer r. For $r \geqslant 0$ let $S_r = (xy)^r S (xy)^{-r}$. Then S_r is a 2^n-dimensional commutative subalgebra of A, which is generated by n elements and satisfies $u^2 \in F$ for every $u \in S_r$. Set $\alpha = x^2 \in F^\times$ and $\beta = y^2 \in F^\times$. Then

$$(xy)^{-r} = (y^{-1}x^{-1})^r = (\beta^{-1}y\alpha^{-1}x)^r = \alpha^{-r}\beta^{-r}(yx)^r.$$

Hence, $S_r = \alpha^{-r}\beta^{-r}(xy)^r S (yx)^r \subseteq \mathrm{Sym}(A,\sigma)$, i.e., S_r is an inseparable subalgebra of (A,σ). However, there exists a finite number of inseparable subalgebras of (A,σ), so $S_r = S_s$ for some non-negative integers r,s with $r > s$. It follows that $S_{r-s} = S_0 = S$. In particular, we have $(xy)^{r-s}y(xy)^{s-r} \in S$ and

$$(xy)^{r-s}y(xy)^{s-r}y = y(xy)^{r-s}y(xy)^{s-r}. \tag{1}$$

Set $\lambda = \alpha^{s-r}\beta^{s-r}$, so that $(xy)^{s-r} = \lambda(yx)^{r-s}$. Replacing in (1) we obtain

$$\lambda(xy)^{r-s}y(yx)^{r-s}y = \lambda y(xy)^{r-s}y(yx)^{r-s}.$$

It follows that $\lambda y^2(xy)^{2(r-s)} = \lambda y^2(yx)^{2(r-s)}$, because $y^2 \in F^\times$. Hence, $(xy)^k = (yx)^k$, where $k = 2(r-s)$. We also have $\sigma((xy)^k) = (yx)^k = (xy)^k$ and $((xy)^k)^2 = (xy)^k(yx)^k \in F^\times$, hence $(xy)^k \in \mathrm{Sym}(A,\sigma)^+$. Finally, we have $(xy)^k x = x(yx)^k = x(xy)^k$, completing the proof. □

Proposition 3.12. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F and let $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$. Then σ is isotropic if and only if $\sigma|_{C_A(x)}$ is isotropic.*

Proof. Observe first that since $x^2 \notin F^2$, $C_A(x)$ is a central simple algebra over $F(x) = F(\sqrt{\alpha})$, where $\alpha = x^2 \in F^\times$. If $\sigma|_{C_A(x)}$ is isotropic, then σ is clearly isotropic. To prove the converse, let S be an inseparable subalgebra of (A,σ). Since σ is isotopic, there exists $y \in S \setminus F$ with $y^2 = 1$. By Lemma 3.11, there is a positive integer k such that $(xy)^k \in \mathrm{Sym}(A,\sigma)^+$. Let r be the minimum positive integer with $(xy)^r \in \mathrm{Sym}(A,\sigma)^+$, hence $(xy)^r = (yx)^r$. We claim that $(xy)^r \neq x^r$. Suppose that $(xy)^r = x^r$. If r is odd, write $r = 2s+1$ for some non-negative integer s. The equality $(xy)^r = x^r$ then implies that $(yx)^s y(xy)^s = x^{2s} = \alpha^s$. As $(xy)^s = \alpha^s(yx)^{-s}$, we get $\alpha^s(yx)^s y(yx)^{-s} = \alpha^s$. Hence, $y = 1 \in F$, which contradicts the assumption. If r is even, write $r = 2s$ for some positive integer s, so that $(xy)^r = x^r = \alpha^s$. Multiplying with $(xy)^{-s}$ we obtain

$$(xy)^s = \alpha^s(xy)^{-s} = \alpha^s\alpha^{-s}(yx)^s = (yx)^s.$$

It follows that $(xy)^s \in \mathrm{Sym}(A,\sigma)^+$, contradicting the minimality of r. This proves the claim. According to Lemma 3.11, we have $(xy)^r \in C_A(x)$. Set $z = (xy)^r + x^r \in C_A(x)$. Then $z \neq 0$ and $\sigma(z)z = \alpha^r + \alpha^r = 0$, i.e., $\sigma|_{C_A(x)}$ is isotropic. □

4 The set of elements represented by σ

Definition 4.1. Let (A,σ) be a totally decomposable algebra with orthogonal involution over F and let $\alpha \in F$. We say that σ *represents* α if there exists a nonzero element $x \in A$ such that $\sigma(x)x + \alpha \in \mathrm{Alt}(A,\sigma)$. The set of all elements in F represented by σ is denoted by $D(A,\sigma)$.

The following result is a restatement of [6, (6.1)].

Lemma 4.2. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F. Then σ is anisotropic if and only if $0 \notin D(A,\sigma)$.*

Lemma 4.3. *Let $\mathfrak{b}$ be an anisotropic bilinear Pfister form over F and let $\alpha \in F$. If $\alpha \notin D(\mathfrak{b})$, then $\mathfrak{b}_{F(\sqrt{\alpha})}$ is anisotropic.*

Proof. Since $\alpha \notin D(\mathfrak{b})$, we have $\alpha \notin F^{\times 2}$, hence $F(\sqrt{\alpha})$ is a field. By [7, (6.1)], $\mathfrak{b} \otimes \langle\langle \alpha \rangle\rangle$ is anisotropic, hence $\mathfrak{b}_{F(\sqrt{\alpha})}$ is also anisotropic by [8, (4.2)]. □

Lemma 4.4. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F. If σ is anisotropic, then $D(A,\sigma) \subseteq D(\mathfrak{Pf}(A,\sigma))$.*

Proof. Let $\alpha \in D(A,\sigma)$. Then there exists $0 \neq x \in A$ such that $\sigma(x)x + \alpha \in \mathrm{Alt}(A,\sigma)$. Suppose that $\alpha \notin D(\mathfrak{Pf}(A,\sigma))$ and set $K = F(\sqrt{\alpha})$. Since $\mathfrak{Pf}(A,\sigma)$ is a Pfister form, [6, (7.5 (1))] implies that $\mathfrak{Pf}(A,\sigma)$ is anisotropic. Hence $\mathfrak{Pf}(A,\sigma)_K$ is also anisotropic by Lemma 4.3. By [6, (7.5 (2))], $(A,\sigma)_K$ is anisotropic. As $\mathrm{Alt}((A,\sigma)_K) = \mathrm{Alt}(A,\sigma)\otimes K$, we have $\sigma_K(x\otimes 1)\cdot(x\otimes 1)+\alpha\otimes 1 \in \mathrm{Alt}((A,\sigma)_K)$. Set $y = x\otimes 1 + 1\otimes\sqrt{\alpha} \in A_K$. Then $y \neq 0$ and

$$\begin{aligned}\sigma_K(y)y &= \sigma_K(x\otimes 1 + 1\otimes\sqrt{\alpha})\cdot(x\otimes 1 + 1\otimes\sqrt{\alpha})\\ &= \sigma_K(x\otimes 1)\cdot(x\otimes 1) + \alpha\otimes 1 + (1\otimes\sqrt{\alpha})(x\otimes 1 + \sigma_K(x\otimes 1)).\end{aligned}$$

We therefore obtain $\sigma_K(y)y \in \mathrm{Alt}((A,\sigma)_K)$, i.e., $0 \in D((A,\sigma)_K)$. This contradicts Lemma 4.2, because $(A,\sigma)_K$ is anisotropic. □

Corollary 4.5. *Let (A,σ) be a totally decomposable algebra with anisotropic orthogonal involution over F. If $\alpha + \sum_{i=1}^m \sigma(x_i)x_i \in \mathrm{Alt}(A,\sigma)$ for some $\alpha \in F$ and $x_1,\cdots,x_m \in A$, then $\alpha \in Q(\mathfrak{Pf}(A,\sigma))$.*

Proof. Set $y = \sum_{i=1}^m x_i \in A$. Then

$$\begin{aligned}\alpha + \sigma(y)y &= \alpha + \sum_{i=1}^m \sigma(x_i)\cdot\sum_{i=1}^m x_i = \alpha + \sum_{i=1}^m \sigma(x_i)x_i + \sum_{i\neq j}\sigma(x_j)x_i\\ &= (\alpha + \sum_{i=1}^m \sigma(x_i)x_i) + \sum_{i<j}(\sigma(x_j)x_i + \sigma(x_i)x_j)\\ &= (\alpha + \sum_{i=1}^m \sigma(x_i)x_i) + \sum_{i<j}(\sigma(\sigma(x_i)x_j) - \sigma(x_i)x_j).\end{aligned}$$

Hence, $\alpha+\sigma(y)y \in \mathrm{Alt}(A,\sigma)$. If $y = 0$, then $\alpha \in \mathrm{Alt}(A,\sigma)$ and the orthogonality of σ implies that $\alpha = 0 \in Q(\mathfrak{Pf}(A,\sigma))$. Otherwise, we have $\alpha \in D(A,\sigma)$ and the result follows from Lemma 4.4. □

Remark 4.6. Let (B,ρ) be a central simple algebra with involution over F and set $(A,\sigma) = (B,\rho)\otimes(M_2(F),t)$. Then every element $x \in A$ can be written as $\left(\begin{smallmatrix} a & b\\ c & d\end{smallmatrix}\right)$, where $a,b,c,d \in B$. The involution σ maps x to $\left(\begin{smallmatrix} \rho(a) & \rho(c)\\ \rho(b) & \rho(d)\end{smallmatrix}\right)$. It follows that

$$\begin{aligned}\mathrm{Alt}(A,\sigma) &= \{\begin{pmatrix} a & b\\ \rho(b) & c\end{pmatrix} \mid a,c \in \mathrm{Alt}(B,\rho) \text{ and } b \in B\},\\ \mathrm{Sym}(A,\sigma) &= \{\begin{pmatrix} a & b\\ \rho(b) & c\end{pmatrix} \mid a,c \in \mathrm{Sym}(B,\rho) \text{ and } b \in B\}.\end{aligned}$$

Lemma 4.7. *If $\alpha I + \sum_{i=1}^m A_i^t A_i \in \mathrm{Alt}(M_2(F),t)$, where $A_1,\cdots,A_m \in M_2(F)$ and $I \in M_2(F)$ is the identity matrix, then $\alpha \in F^2$.*

Proof. Write $A_i = \left(\begin{smallmatrix} a_i & b_i\\ c_i & d_i\end{smallmatrix}\right)$ for some $a_i,b_i,c_i,d_i \in F$, so that

$$A_i^t A_i = \begin{pmatrix} a_i^2 + c_i^2 & a_ib_i + c_id_i\\ a_ib_i + c_id_i & b_i^2 + d_i^2\end{pmatrix}.$$

Since $\mathrm{Alt}(M_2(F),t)$ is the set of symmetric matrices with zero diagonal, the assumption implies that $\alpha + \sum_{i=1}^m a_i^2 + c_i^2 = 0$, hence $\alpha \in F^2$. □

Lemma 4.8. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F. If $\alpha+\sum_{i=1}^{m}\sigma(x_i)x_i \in \mathrm{Alt}(A,\sigma)$ for some $\alpha \in F$ and $x_1,\cdots,x_m \in A$, then $\alpha \in Q(\mathfrak{Pf}(A,\sigma))$.*

Proof. Let $\deg_F A = 2^n$. We use induction on n. If σ is anisotropic, the conclusion follows from Corollary 4.5. Suppose that σ is isotropic. If $n=1$, then $(A,\sigma) \simeq (M_2(F),t)$ and $\mathfrak{Pf}(A,\sigma) \simeq \langle\langle 1\rangle\rangle$ and the result follows from Lemma 4.7. Let $n \geqslant 2$. Since $\mathfrak{i}(A,\sigma) > 0$, we may identify $(A,\sigma) = (B,\rho)\otimes(M_2(F),t)$, where (B,ρ) is a totally decomposable algebra with orthogonal involution over F. Then $\mathfrak{Pf}(A,\sigma) \simeq \mathfrak{Pf}(B,\rho)\otimes\langle\langle 1\rangle\rangle$, which implies that $Q(\mathfrak{Pf}(A,\sigma)) = Q(\mathfrak{Pf}(B,\rho))$. Write $x_i = \left(\begin{smallmatrix} a_i & b_i \\ c_i & d_i \end{smallmatrix}\right)$, where $a_i,b_i,c_i,d_i \in B$. Using Remark 4.6 we get

$$\sigma(x_i)x_i = \begin{pmatrix} \rho(a_i) & \rho(c_i) \\ \rho(b_i) & \rho(d_i) \end{pmatrix} \cdot \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix}.$$

Hence the $(1,1)$-entry of $\sum_{i=1}^{m}\sigma(x_i)x_i$ is $\sum_{i=1}^{m}(\rho(a_i)a_i + \rho(c_i)c_i)$. Since $\alpha + \sum_{i=1}^{m}\sigma(x_i)x_i \in \mathrm{Alt}(A,\sigma)$, Remark 4.6 implies that

$$\alpha + \sum_{i=1}^{m}(\rho(a_i)a_i + \rho(c_i)c_i) \in \mathrm{Alt}(B,\rho).$$

By induction hypothesis we get $\alpha \in Q(\mathfrak{Pf}(B,\rho)) = Q(\mathfrak{Pf}(A,\sigma))$. □

Corollary 4.9. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F. If $x \in \mathrm{Sym}(A,\sigma)^+$, then $x^2 \in Q(\mathfrak{Pf}(A,\sigma))$.*

Proof. Set $\alpha = x^2 \in F$. Then $\alpha + \sigma(x)x = 0 \in \mathrm{Alt}(A,\sigma)$ and the result follows from Lemma 4.8. □

Theorem 4.10. *If (A,σ) is a totally decomposable algebra with orthogonal involution over F, then $D(A,\sigma) = D(\mathfrak{Pf}(A,\sigma)) = \{x^2 \mid 0 \neq x \in S\}$, where S is any inseparable subalgebra of (A,σ).*

Proof. If $\alpha \in D(\mathfrak{Pf}(A,\sigma))$, then there exists $0 \neq x \in S$ such that $x^2 = \mathfrak{Pf}(A,\sigma)(x,x) = \alpha$. Hence, $\sigma(x)x + \alpha = 0 \in \mathrm{Alt}(A,\sigma)$, i.e., $\alpha \in D(A,\sigma)$. Conversely, let $\alpha \in D(A,\sigma)$. If $\alpha = 0$, then σ is isotropic by Lemma 4.2. Hence, $\mathfrak{Pf}(A,\sigma)$ is isotopic by [6, (7.5) and (6.2)], i.e, $0 \in D(\mathfrak{Pf}(A,\sigma))$. Otherwise, as $Q(\mathfrak{Pf}(A,\sigma)) = D(\mathfrak{Pf}(A,\sigma)) \cup \{0\}$, the result follows from Lemma 4.8. The equality $D(\mathfrak{Pf}(A,\sigma)) = \{x^2 \mid 0 \neq x \in S\}$ follows from the relation $\mathfrak{Pf}(A,\sigma)(x,x) = x^2$ for $x \in S$. □

Lemma 4.11. *Let $\mathfrak{b}$ be a bilinear n-fold Pfister form over F. If $\alpha \in Q(\mathfrak{b}) \setminus F^2$, then $\mathfrak{i}(\mathfrak{b}_{F(\sqrt{\alpha})}) = \mathfrak{i}(\mathfrak{b}) + 1$.*

Proof. Set $K = F(\sqrt{\alpha})$ and $r = \mathfrak{i}(\mathfrak{b})$. As $Q(\langle\langle 1\rangle\rangle) = F^2$ and $\alpha \in Q(\mathfrak{b}) \setminus F^2$ we have $\mathfrak{b} \not\simeq \langle\langle 1\rangle\rangle^n$, i.e., $r < n$. Write $\mathfrak{b} \simeq \langle\langle 1\rangle\rangle^r \otimes \mathfrak{b}'$ for some anisotropic bilinear Pfister form $\mathfrak{b}'$ over F. Since $Q(\mathfrak{b}) = Q(\mathfrak{b}')$, we have $\alpha \in Q(\mathfrak{b}')$. Hence, the pure subform of $\mathfrak{b}'$ represents $\alpha + \lambda^2$ for some $\lambda \in F$. By [1, A.2], there exist $\alpha_2,\cdots,\alpha_s \in F$ such that $\mathfrak{b}' \simeq \langle\langle \alpha+\lambda^2, \alpha_2,\cdots,\alpha_s\rangle\rangle$. Note that we have $\alpha+\lambda^2 \in K^{\times 2}$, hence $\mathfrak{b}'_K \simeq \langle\langle 1, \alpha_2,\cdots,\alpha_s\rangle\rangle_K$. Since $\mathfrak{b}' = \langle\langle \alpha+\lambda^2\rangle\rangle\otimes\langle\langle \alpha_2,\cdots,\alpha_s\rangle\rangle$ is anisotropic and $K = F(\sqrt{\alpha+\lambda^2})$, by [8, (4.2)] the form $\langle\langle \alpha_2,\cdots,\alpha_s\rangle\rangle_K$ is anisotropic. Hence $\mathfrak{i}(\mathfrak{b}'_K) = 1$, which implies that $\mathfrak{i}(\mathfrak{b}_K) = r+1 = \mathfrak{i}(\mathfrak{b})+1$. □

Corollary 4.12. *Let (A,σ) be a totally decomposable algebra with orthogonal involution over F. If $x \in \mathrm{Sym}(A,\sigma)^+$ with $\alpha := x^2 \notin F^{\times 2}$, then $\mathfrak{i}((A,\sigma)_{F(\sqrt{\alpha})}) = \mathfrak{i}(A,\sigma)+1$. In particular, $(A,\sigma) \not\simeq (M_{2^n}(F),t)$.*

Proof. By Corollary 4.9, we have $\alpha \in Q(\mathfrak{Pf}(A,\sigma))$. Hence, the result follows from Lemma 4.11. □

5 Stable quaternion subalgebras

In this section we study some conditions under which a symmetric square-central element of a totally decomposable algebra with orthogonal involution is contained in a stable quaternion subalgebra. We start with anisotropic involutions.

Theorem 5.1. *Let (A,σ) be a totally decomposable algebra with anisotropic orthogonal involution over F. Then every $x \in \mathrm{Sym}(A,\sigma)^+$ is contained in a σ-invariant quaternion subalgebra of A.*

Proof. Since σ is anisotropic, Theorem 3.5 shows that x is contained in the unique inseparable subalgebra of (A,σ). If $x^2 = \lambda^2$ for some $\lambda \in F$, then $(x+\lambda)^2 = 0$. Hence, $x = \lambda$ by [6, (6.1)] and the result is trivial. Otherwise, $x^2 \notin F^2$ and the conclusion follows from [14, (6.3 (ii))]. □

We next consider algebras of degree 4 and 8.

Proposition 5.2. *Let (A,σ) be a totally decomposable algebra of degree 4 with orthogonal involution over F. If $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$, then x is contained in a σ-invariant quaternion subalgebra of A.*

Proof. By Corollary 4.12 we have either $\mathfrak{i}(A,\sigma) = 0$ or $\mathfrak{i}(A,\sigma) = 1$. In the first case, the result follows from Theorem 5.1. Suppose that $\mathfrak{i}(A,\sigma) = 1$. Set $C = C_A(x)$ and $K = F(x) = F(\sqrt{\alpha})$. By Proposition 3.12, $(C,\sigma|_C)$ is isotropic, i.e., $(C,\sigma|_C) \simeq (M_2(K),t) \simeq (M_2(F),t)_K$. Hence, the algebra $M_2(F)$ may be identified with a subalgebra of $C \subseteq A$. Let $Q = C_A(M_2(F))$. Then $(Q,\sigma|_Q)$ is a σ-invariant quaternion subalgebra of A containing x. □

Lemma 5.3. *Let $\mathfrak{b}$ be a 2-dimensional anisotropic symmetric bilinear form over F and let $K = F(\sqrt{\alpha})$ for some $\alpha \in F^\times \setminus F^{\times 2}$. If $\mathfrak{b}_K$ is isotropic, then $\mathfrak{b} \simeq \langle a, sa\rangle$ for some $a \in F^\times$ and $s \in K^{\times 2}$.*

Proof. The result follows from [12, (3.5)]. □

We recall that a symmetric bilinear space $(V,\mathfrak{b})$ over F is called *metabolic* if there exists a subspace W of V with $\dim_F W = \frac{1}{2}\dim_F V$ such that $\mathfrak{b}|_{W\times W} = 0$.

Lemma 5.4. *Let $\mathfrak{b}$ be a 4-dimensional isotropic symmetric bilinear form over F. Let $K = F(\sqrt{\alpha})$ for some $\alpha \in F^\times \setminus F^{\times 2}$. If $\mathfrak{b} \otimes \langle\langle \alpha\rangle\rangle$ is a Pfister form, then $\mathfrak{b}_K$ is similar to a Pfister form.*

Proof. Since $\mathfrak{b} \otimes \langle\langle \alpha\rangle\rangle$ is a Pfister form, the form $\mathfrak{b}$ is not alternating. If $\mathfrak{b}$ is metabolic, then by [7, (1.24) and (1.22 (3))] either $\mathfrak{b} \simeq \langle a,a,b,b\rangle$ or $\mathfrak{b} \simeq \langle a,a\rangle \perp \mathbb{H}$, where $a,b \in F^\times$ and $\mathbb{H}$ is the hyperbolic plane. In the first case $\mathfrak{b}$ is similar to $\langle 1,1,ab,ab\rangle = \langle\langle 1,ab\rangle\rangle$. In the second case, using the isometry

$\langle a,a,a\rangle \simeq \langle a\rangle \perp \mathbb{H}$ in [7, (1.16)], we get $\mathfrak{b} \simeq \langle a,a,a,a\rangle$. Hence, $\mathfrak{b}$ is similar to $\langle\langle 1,1\rangle\rangle$.

Suppose that $\mathfrak{b}$ is not metabolic. By Witt decomposition theorem [7, (1.27)] and the isometry $\langle a,a,a\rangle \simeq \langle a\rangle \perp \mathbb{H}$ for $a \in F^\times$, the form $\mathfrak{b}$ is similar to $\mathfrak{b}' \perp \langle 1,1\rangle$ for some 2-dimensional anisotropic bilinear form $\mathfrak{b}'$ over F. Since $\mathfrak{b} \otimes \langle 1,\alpha\rangle$ is an isotropic Pfister form, it is metabolic by [7, (6.3)]. Hence, $\mathfrak{b}' \otimes \langle 1,\alpha\rangle$ is also metabolic. By [8, (4.2)] the form $\mathfrak{b}'_K$ is metabolic. Using Lemma 5.3, one can write $\mathfrak{b}' \simeq \langle a, sa\rangle$, where $a \in F^\times$ and $s \in K^{\times 2}$. The form $\mathfrak{b}_K$ is therefore similar to $\langle 1,1,a,sa\rangle_K \simeq \langle 1,1,a,a\rangle_K \simeq \langle\langle 1,a\rangle\rangle_K$, proving the result □

Lemma 5.5. *Let (A,σ) be a central simple algebra of degree 4 with orthogonal involution over F and let K/F be a separable quadratic extension. If $(A,\sigma)_K$ is totally decomposable, then (A,σ) is also totally decomposable.*

Proof. The result follows from [11, (7.3)] and the relation $K^{\times 2} \cap F^\times = F^{\times 2}$. □

Lemma 5.6. *Let (A,σ) be a totally decomposable algebra of degree 2^n with orthogonal involution over F. Let $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$ and set $C = C_A(x)$. If $(C,\sigma|_C)$ is totally decomposable and x is contained in a σ-invariant quaternion subalgebra Q, then there exists an inseparable subalgebra of (A,σ) containing x.*

Proof. Let $\alpha = x^2 \in F^\times \setminus F^{\times 2}$, $K = F(x) = F(\sqrt{\alpha})$, $B = C_A(Q)$ and $\rho = \sigma|_B$. Then there exists a natural isomorphism of K-algebras with involution

$$f : (B,\rho)_K \simeq (C,\sigma|_C).$$

In particular, $(B,\rho)_K$ is totally decomposable. By [15, (3.6)], there exists a totally decomposable algebra of degree 2^{n-1} with orthogonal involution (B',ρ') over F such that $(B,\rho)_K \simeq (B',\rho')_K$. Let $S = F[v_1,\cdots,v_{n-1}]$ be an inseparable subalgebra of (B',ρ'). The algebra $K[v_1,\cdots,v_{n-1}] \simeq S_K$ may be identified with an inseparable subalgebra of $(B,\rho)_K$. Set $v'_i = f(v_i) \in C$, $i = 1,\cdots,n-1$. Then $S' := K[v'_1,\cdots,v'_{n-1}]$ is an inseparable subalgebra of $(C,\sigma|_C)$ satisfying $u^2 \in F$ for $u \in S'$. Since $(C,\sigma|_C) \subseteq (A,\sigma)$, we have $S' \subseteq \mathrm{Sym}(A,\sigma)^+$. By [14, (3.11)], S' is a Frobenius subalgebra of A, because $\dim_F S' = 2^n$ and S' is generated, as an F-algebra, by $x, v'_1,\cdots,v'_{n-1}$. It follows from [9, (2.2.3)] that $C_A(S') = S'$. Hence, S' is an inseparable subalgebra of (A,σ) containing x. □

Proposition 5.7. *Let (A,σ) be a totally decomposable algebra of degree 8 over F. For an element $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$, the following conditions are equivalent:*

(1) There exists a σ-invariant quaternion subalgebra of A containing x.

(2) There exists an inseparable subalgebra S of (A,σ) such that $x \in S$.

Proof. If $\mathrm{i}(A,\sigma) = 0$, by Theorem 5.1 and Theorem 3.5 both conditions are satisfied. Let $\mathrm{i}(A,\sigma) > 0$. Then $(A,\sigma) \simeq (M_2(F),t) \otimes (Q_1,\sigma_1) \otimes (Q_2,\sigma_2)$, where (Q_i,σ_i), $i = 1,2$, is a quaternion algebra with orthogonal involution over F. Suppose first that x is contained in a σ-invariant quaternion subalgebra Q_3 of

A. Set $B = C_A(Q_3)$, $\sigma_3 = \sigma|_{Q_3}$ and $\rho = \sigma|_B$, so that $(A,\sigma) \simeq (Q_3,\sigma_3) \otimes (B,\rho)$. It follows that

$$(Q_3,\sigma_3) \otimes (B,\rho) \simeq (M_2(F),t) \otimes (Q_1,\sigma_1) \otimes (Q_2,\sigma_2). \tag{2}$$

Let $C = C_A(x)$ and $K = F(x) = F(\sqrt{\alpha})$, where $\alpha = x^2 \in F^\times \setminus F^{\times 2}$. Then $(C,\sigma|_C) \simeq (B,\rho)_K$ as K-algebras. By Proposition 3.12, $(C,\sigma|_C)$ is isotropic, hence $(B,\rho)_K$ is also isotropic. We claim that $(B,\rho)_K$ is totally decomposable. The result then follows from Lemma 5.6.

By Lemma 3.8, for $i = 1,2,3$ there exists a quaternion basis $(1,u_i,v_i,w_i)$ of Q_i such that $u_i \in \mathrm{Sym}(Q_i,\sigma_i)$. Let $\beta_i = u_i^2 + u_i \in F$. For $i = 0,1,2,3$, define a field L_i inductively as follows: set $L_0 = F$. For $i \geqslant 1$ set $L_i = L_{i-1}(u_i)$ if $\beta_i \notin \wp(L_{i-1}) := \{y^2 + y \mid y \in L_{i-1}\}$ and $L_i = L_{i-1}$ otherwise. In other words, either $L_i = L_{i-1}$ or L_i/L_{i-1} is a separable quadratic extension. Note that we have $L_i^{\times 2} \cap F^\times = F^{\times 2}$, hence either $L_i(\sqrt{\alpha}) = L_{i-1}(\sqrt{\alpha})$ or $L_i(\sqrt{\alpha})/L_{i-1}(\sqrt{\alpha})$ is a separable quadratic extension. We show that $\rho_{L_3(\sqrt{\alpha})}$ is totally decomposable, which implies that $\rho_{L_i(\sqrt{\alpha})}$ is totally decomposable for $i = 0,1,2$, thanks to Lemma 5.5. In particular, $\rho_K = \rho_{F(\sqrt{\alpha})}$ is also totally decomposable, as required.

Set $L = L_3$. For $i = 1,2,3$, the algebra Q_{iL} splits. Hence we may identify $(Q_i,\sigma_i)_L = (M_2(L),\tau_i)$, where τ_i is an orthogonal involution on $M_2(L)$. Using (2) we obtain

$$(M_2(L),\tau_3) \otimes (B,\rho)_L \simeq (M_2(L),t) \otimes (M_2(L),\tau_1) \otimes (M_2(L),\tau_2). \tag{3}$$

Hence, B_L splits and we may identify $(B,\rho)_L = \mathrm{Ad}(\mathfrak{b})$ for some symmetric bilinear form $\mathfrak{b}$ over L. As $(B,\rho)_L$ is isotropic, the form $\mathfrak{b}$ is isotropic by [6, (4.3)]. Since $x \in \mathrm{Sym}(Q_3,\sigma_3)^+ \setminus F$, by [16, (5.4)] there exists $\lambda \in F$ for which $0 \neq x + \lambda \in \mathrm{Alt}(Q_3,\sigma_3)$. Hence, $\operatorname{disc}\sigma_3 = (\alpha + \lambda^2)F^{\times 2}$, which implies that

$$(M_2(L),\tau_3) \simeq (Q_3,\sigma_3)_L \simeq \mathrm{Ad}(\langle\!\langle \alpha + \lambda^2 \rangle\!\rangle_L), \tag{4}$$

by [10, (7.4)]. The right side of (3) is the adjoint involution of a 3-fold bilinear Pfister form over L. Hence, using (4) and [10, (4.2)] it follows that $\mathfrak{b} \otimes \langle\!\langle \alpha + \lambda^2 \rangle\!\rangle$ is similar to a Pfister form over L. By Lemma 5.4, $\mathfrak{b}_{L(\sqrt{\alpha})}$ is similar to a Pfister form. Hence, $\rho_{L(\sqrt{\alpha})}$ is totally decomposable, as required. This proves the implication (1) $\Rightarrow$ (2). The converse follows from [14, (6.3 (ii))]. □

Lemma 5.8. *If $\mathfrak{b}$ is a 2^n-dimensional symmetric bilinear form over F, then $\mathrm{Ad}(\mathfrak{b}) \simeq (M_{2^n}(F),t)$ if and only if $\mathfrak{b} \simeq \lambda \cdot \langle\!\langle 1 \rangle\!\rangle^n$ for some $\lambda \in F^\times$.*

Proof. The result follows from [14, (5.7)], [6, (7.5 (3))] and [10, (4.2)]. □

Lemma 5.9. *Let (A,σ) be a central simple algebra with orthogonal involution over F. If $(A,\sigma) \otimes (M_2(F),\tau) \simeq (M_{2^n}(F),t)$ for some orthogonal involution τ on $M_2(F)$, then $(A,\sigma) \simeq (M_{2^{n-1}}(F),t)$.*

Proof. Observe first that A splits, hence we may identify $(A,\sigma) = \mathrm{Ad}(\mathfrak{b})$ for some symmetric bilinear form $\mathfrak{b}$ over F. By [10, (7.3 (3)) and (7.4)], we have $(M_2(F),\tau) \simeq \mathrm{Ad}(\langle\!\langle \alpha \rangle\!\rangle)$, where $\alpha \in F^\times$ is a representative of the class $\operatorname{disc}\tau \in F^\times/F^{\times 2}$. Also, $(M_{2^n}(F),t) \simeq \mathrm{Ad}(\langle\!\langle 1 \rangle\!\rangle^n)$ by Lemma 5.8. The assumption implies that $\mathrm{Ad}(\mathfrak{b} \otimes \langle\!\langle \alpha \rangle\!\rangle) \simeq \mathrm{Ad}(\langle\!\langle 1 \rangle\!\rangle^n)$. Hence, the forms $\mathfrak{b} \otimes \langle\!\langle \alpha \rangle\!\rangle$ and $\langle\!\langle 1 \rangle\!\rangle^n$ are similar by [10, (4.2)], i.e., there exists $\lambda \in F^\times$ for which $\mathfrak{b} \otimes \langle\!\langle \alpha \rangle\!\rangle \simeq \lambda \cdot \langle\!\langle 1 \rangle\!\rangle^n$. As $Q(\langle\!\langle 1 \rangle\!\rangle^n) = F^2$, we obtain $Q(\mathfrak{b}) \subseteq \lambda \cdot F^2$, i.e., $\mathfrak{b}$ is similar to $\langle\!\langle 1 \rangle\!\rangle^{n-1}$. Using Lemma 5.8 we get $(A,\sigma) \simeq (M_{2^{n-1}}(F),t)$. □

Theorem 5.10. *Let (A,σ) be a totally decomposable algebra of degree 2^n with orthogonal involution over F and let $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$. If $\mathfrak{i}(A,\sigma) = n-1$, then the following statements are equivalent:*

(1) There exists a σ-invariant quaternion subalgebra Q of A containing x.

(2) There exists an inseparable subalgebra S of (A,σ) such that $x \in S$.

Proof. The implication (2) $\Rightarrow$ (1) follows from [14, (6.3 (ii))]. To prove the converse, let $C = C_A(x)$. In view of Lemma 5.6, it suffices to show that $(C, \sigma|_C)$ is totally decomposable. Let $\tau = \sigma|_Q$, $B = C_A(Q)$ and $\rho = \sigma|_B$. Then $(A,\sigma) \simeq (B,\rho) \otimes (Q,\tau)$. Set $K = F(x) = F(\sqrt{\alpha})$, where $\alpha = x^2 \in F^\times \setminus F^{\times 2}$. We have $(C,\sigma|_C) \simeq_K (B,\rho)_K$. Hence, it is enough to show that $(B,\rho)_K$ is totally decomposable. By Corollary 4.12 we have $\mathfrak{i}(A,\sigma)_K = n$, so $(A,\sigma)_K \simeq (M_{2^n}(K), t)$. It follows that $(B,\rho)_K \otimes_K (Q,\tau)_K \simeq (M_{2^n}(K),t)$. Since $x \in Q$ and $x^2 = \alpha \in K^2$, the algebra Q_K splits. Hence, using Lemma 5.9 we get $(B,\rho)_K \simeq_K (M_{2^{n-1}}(K),t)$. In particular, $(B,\rho)_K$ is totally decomposable, proving the result. □

6 Examples for isotropic involutions

In this section we show that the criteria obtained in §5 do not necessarily apply to arbitrary involutions.

Lemma 6.1. *Let (A,σ) be a totally decomposable algebra of degree 2^n with orthogonal involution over F. If $n \geqslant 2$ and $(A,\sigma) \not\simeq (M_{2^n}(F),t)$, then there exist an element $w \in \mathrm{Sym}(A,\sigma) \setminus (\mathrm{Alt}(A,\sigma) \oplus F)$ and a unit $u \in \mathrm{Alt}(A,\sigma)$ such that $u^2 \in F^\times \setminus F^{\times 2}$ and $uw = wu$.*

Proof. Let $(A,\sigma) \simeq \bigotimes_{i=1}^n (Q_i,\sigma_i)$ be a decomposition of (A,σ). As $(A,\sigma) \not\simeq (M_{2^n}(F),t)$, (by re-indexing) we may assume that $(Q_1,\sigma_1) \not\simeq (M_2(F),t)$. Let $u \in \mathrm{Alt}(Q_1,\sigma_1)$ be a unit, so that $u^2 \in F^\times$. Note that $u^2 \notin F^{\times 2}$, since otherwise Q_1 splits and $\mathrm{disc}\,\sigma_1$ is trivial. As $\mathrm{disc}\,t$ is also trivial (see [10, p. 82]), we get $(Q_1,\sigma_1) \simeq (M_2(F),t)$ by [10, (7.4)], contradicting the assumption. Hence, $u^2 \in F^\times \setminus F^{\times 2}$. By [10, (2.6)] we have $\dim_F \mathrm{Sym}(Q_2,\sigma_2) = 3$ and $\dim_F \mathrm{Alt}(Q_2,\sigma_2) = 1$. Hence there exists an element $w \in \mathrm{Sym}(Q_2,\sigma_2) \setminus (\mathrm{Alt}(Q_2,\sigma_2) \oplus F)$. Clearly, we have $uw = wu$. The elements u and w may be identified with elements of A, so that $w \in \mathrm{Sym}(A,\sigma)$ and $u \in \mathrm{Alt}(A,\sigma)$. Since $\alpha + w \notin \mathrm{Alt}(Q_2,\sigma_2)$ for every $\alpha \in F$, by [13, (3.5)] we have $\alpha + w \notin \mathrm{Alt}(A,\sigma)$ for all $\alpha \in F$, i.e., $w \in \mathrm{Sym}(A,\sigma) \setminus (\mathrm{Alt}(A,\sigma) \oplus F)$, as required. □

The next result shows that Theorem 5.1 does not hold for isotropic involutions of degree $\geqslant 8$ (see also Proposition 5.2).

Proposition 6.2. *Let (A,σ) be a totally decomposable algebra of degree 2^n with isotropic orthogonal involution over F. If $n \geqslant 3$ and $(A,\sigma) \not\simeq (M_{2^n}(F),t)$, then there exists an element $x \in \mathrm{Sym}(A,\sigma)^+$ with $x^2 \notin F^2$ which is not contained in any σ-invariant quaternion subalgebra of A.*

Proof. Since $\mathfrak{i}(A,\sigma) > 0$, we may identify $(A,\sigma) = (B,\rho) \otimes (M_2(F),t)$, where (B,ρ) is a totally decomposable algebra with orthogonal involution over F. The assumptions $n \geqslant 3$ and $(A,\sigma) \not\simeq (M_{2^n}(F),t)$ imply that $\deg_F B \geqslant 4$ and

$(B,\rho) \not\simeq (M_{2^{n-1}}(F), t)$. By Lemma 6.1, there exists an element $w \in \mathrm{Sym}(B,\rho) \setminus (\mathrm{Alt}(B,\rho) \oplus F)$ and a unit $u \in \mathrm{Alt}(B,\rho)$ for which $uw = wu$. Set

$$x = \begin{pmatrix} w & w+u \\ w+u & w \end{pmatrix} \in A.$$

By Remark 4.6 we have $x \in \mathrm{Sym}(A,\sigma) \setminus (\mathrm{Alt}(A,\sigma) \oplus F)$. As $u^2 \in F^\times \setminus F^{\times 2}$ we have $x^2 \in F^\times \setminus F^{\times 2}$. By [16, (5.4)], the element x is not contained in any σ-invariant quaternion subalgebra of A, because $x + \alpha \notin \mathrm{Alt}(A,\sigma)$ for every $\alpha \in F$. □

We conclude by showing that the implication (1) ⇒ (2) in Proposition 5.10 and Proposition 5.7 does not hold for arbitrary involutions. We use the ideas of [6, (9.4)]. Recall that the *canonical* involution γ on a quaternion F-algebra Q is defined as $\gamma(x) = x - \mathrm{Trd}_Q(x)$ for $x \in Q$, where $\mathrm{Trd}_Q(x)$ is the reduced trace of x in Q. Also, for a division algebra with involution (D,θ) over F and $\alpha_1, \cdots, \alpha_n \in D^\times \cap \mathrm{Sym}(D,\theta)$, the diagonal hermitian form h on D^n defined by $h(x,y) = \sum_{i=1}^n \theta(x_i)\alpha_i y_i$ is denoted by $\langle \alpha_1, \cdots, \alpha_n \rangle_\theta$.

Example 6.3. Let $F \neq F^2$ and let $K = F(X,Y,Z)$, where X, Y and Z are indeterminates. Let $Q = [X,Y)_K$ and let γ be the canonical involution on Q. By [6, (9.3)], Q is a division algebra over K. Choose an element $s \in \mathrm{Sym}(Q,\gamma)$ with $s^2 = Y$. Let ψ be the diagonal hermitian form $\langle 1, Z, s, s\rangle_\gamma$ over (Q,γ) and set $(B,\rho) = \mathrm{Ad}(\psi)$. By [6, (9.4)], (B,ρ) is not totally decomposable, but $(B,\rho)_L$ is totally decomposable for every splitting field L of A.

Now, choose $\alpha \in F^\times \setminus F^{\times 2}$ and let $Q' = [X,\alpha)_K$ with a quaternion basis $(1,u,v,w)$. Let τ be the involution on Q' induced by $\tau(u) = u$ and $\tau(v) = v$. Then τ is an orthogonal involution and $v = \tau(uv) - uv \in \mathrm{Alt}(Q',\tau)$. Set $(A,\sigma) = (B,\rho) \otimes_K (Q',\tau)$. Then (A,σ) is a central simple algebra with orthogonal involution over K. We claim that (A,σ) is totally decomposable. Let $L = K(u) \subseteq Q'$ and set $C = C_A(1 \otimes u)$. Then L/K is a separable quadratic extension and

$$(C, \sigma|_C) \simeq_L (B,\rho)_L, \tag{5}$$

is a central simple L-algebra with orthogonal involution. Since $u^2 + u = X$, $Q_L \simeq [X,Y)_L$ splits, which implies that B_L is also split. It follows that $(B,\rho)_L$ is totally decomposable, i.e., $(C,\sigma|_C)$ is totally decomposable by (5). Using [15, (7.3)] and the isomorphism (5), one can find a totally decomposable algebra with orthogonal involution (C',σ') over K such that $(C,\sigma|_C) \simeq (C',\sigma')_L$. As $C \subseteq A$, the algebra C' may be identified with a subalgebra of A. Let $Q'' = C_A(C')$. Then Q'' is a quaternion K-subalgebra of A and $(A,\sigma) \simeq_K (C',\sigma') \otimes_K (Q'', \sigma|_{Q''})$ is totally decomposable, proving the claim.

The element $1 \otimes v \in \mathrm{Alt}(A,\sigma)^+$ is contained in the copy of Q' in A, which is a σ-invariant quaternion subalgebra of A. Note that $(C_A(1 \otimes v), \sigma|_{C_A(1\otimes v)}) \simeq (B,\rho)_{K(v)}$ as $K(v)$-algebras. We show that $(B,\rho)_{K(v)}$ is not totally decomposable, which implies that $1 \otimes v$ is not contained in any inseparable subalgebra of (A,σ), thanks to [14, (6.3)]. Since $v^2 = \alpha \in F^\times \setminus F^{\times 2}$, we have $K(v) \simeq F(\sqrt{\alpha})(X,Y,Z)$. Hence, $Q_{K(v)}$ is still a division algebra by [6, (9.3)]. Using [6, (9.4)] it follows that $(B,\rho)_{K(v)}$ is not totally decomposable.

References

[1] J. Arason, R. Baeza, Relations in I^n and I^nW_q in characteristic 2. *J. Algebra* **314** (2007), no. 2, 895–911.

[2] D. Barry, Decomposable and indecomposable algebras of degree 8 and exponent 2 (with an appendix by A. S. Merkurjev). *Math. Z.* **276** (2014), no. 3-4, 1113–1132.

[3] D. Barry, Power-central elements in tensor products of symbol algebras. *Comm. Algebra* **44** (2016), no. 9, 3767–3787.

[4] D. Barry, A. Chapman, Square-central and Artin-Schreier elements in division algebras. *Arch. Math. (Basel)* **104** (2015), no. 6, 513–521.

[5] A. Dolphin, Decomposition of algebras with involution in characteristic 2. *J. Pure Appl. Algebra* **217** (2013), no. 9, 1620–1633.

[6] A. Dolphin, Orthogonal Pfister involutions in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 10, 1900–1915.

[7] R. Elman, N. Karpenko, A. Merkurjev, *The algebraic and geometric theory of quadratic forms.* American Mathematical Society Colloquium Publications, 56. American Mathematical Society, Providence, RI, 2008.

[8] D. Hoffmann, Witt kernels of bilinear forms for algebraic extensions in characteristic 2. *Proc. Amer. Math. Soc.*, **134** (2006), no. 3, 645–652.

[9] N. Jacobson, *Finite-dimensional division algebras over fields.* Springer-Verlag, Berlin, 1996.

[10] M.-A. Knus, A. S. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions.* American Mathematical Society Colloquium Publications, 44. American Mathematical Society, Providence, RI, 1998.

[11] M.-A. Knus, R. Parimala, R. Sridharan, Involutions on rank 16 central simple algebras. *J. Indian Math. Soc. (N.S.)* **57** (1991), no. 1–4, 143–151.

[12] A. Laghribi, P. Mammone. Hyper-isotropy of bilinear forms in characteristic 2. *Contemporary Mathematics*, **493** (2009), 249-269.

[13] M. G. Mahmoudi, A.-H. Nokhodkar, On split products of quaternion algebras with involution in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 4, 731–734.

[14] M. G. Mahmoudi, A.-H. Nokhodkar, On totally decomposable algebras with involution in characteristic two. *J. Algebra* **451** (2016), 208–231.

[15] A.-H. Nokhodkar, Quadratic descent of totally decomposable orthogonal involutions in characteristic two. *J. Pure Appl. Algebra* **221** (2017), no. 4, 948–959.

[16] A.-H. Nokhodkar, On the decomposition of metabolic involutions. *J. Algebra Appl.* (to appear).

[17] A.-H. Nokhodkar, On decomposable biquaternion algebras with involution of orthogonal type. LAG preprint server, http://www.math.uni-bielefeld.de/LAG/man/558.html.

[18] A. Quéguiner-Mathieu, J.-P. Tignol, Discriminant and Clifford algebras. *Math. Z.* **240** (2002), no. 2, 345–384.

[19] L. H. Rowen, Central simple algebras. *Israel J. Math.* **29** (1978), 285–301.

A.-H. Nokhodkar, anokhodkar@yahoo.com,
Department of Pure Mathematics, Faculty of Science, University of Kashan, P. O. Box 87317-51167, Kashan, Iran.